

General Information for Individuals with a *MC1R* Risk Allele or Mutation

Cancer Risks and General Management Considerations

The cancer risks for individuals who have an *MC1R* mutation are still being understood. However, it has been suggested that individuals with one mutation in the *MC1R* gene may be at an increased chance to develop skin cancer, particularly melanoma. The exact lifetime risk for individuals with a mutation in the *MC1R* gene have not been established, but may indicated a slightly increased risk for melanoma (2.6%-11.8%)¹⁻³ and may confer a slightly increased risk for basal cell carcinoma. Associations with other cancers have not been reported for individuals with a *MC1R* risk allele or mutation at this time.

No consensus management recommendations or guidelines have been published for individuals with a mutation in the *MC1R* gene at this time. Individuals may consider a formal dermatological evaluation based on the increased chance to develop skin cancer.

Implications for Family Members/Reproductive Considerations

- First-degree relatives (i.e., parents, siblings, and children) have a 50% chance to have the familial *MC1R* mutation. Second-degree relatives (i.e., nieces/nephews, aunts/uncles, and grandparents) have a 25% chance to have the familial mutation.
- Individuals with two *MC1R* mutations (one inherited from each parent) have a chance for red hair and paler skin.
- Family members may pursue genetic counseling to clarify their risks. Family members can visit www.FindAGeneticCounselor.com to find genetic services near them.

References

1. Raimondi S, Sera F, Gandini S, et al. 2008. *MC1R* Variants, Melanoma, and Red Hair Color Phenotype: a Meta-Analysis. *Int J Cancer*. 15;122(12):2753-60.
2. Landi Mt, Kanetsky PA, Tsang S, et al. 2005. *MC1R*, *ASIP*, and DNA Repair in Sporadic and Familial Melanoma in a Mediterranean Population. *J Natl Cancer Inst*. 97(13):998-1007.
3. Williams PF, Olsen CM, Hayward NK, Whiteman DC. 2011. Melanocortin 1 Receptor and Risk of Cutaneous Melanoma: a Meta-analysis and Estimate of Population Burden. *Int J Cancer*. 129(7);1730-40.
4. King RA, Willaert RK, Schmidt RM, et al. 2003. *MCR1* Mutations Modify the Classic Phenotype of Oculocutaneous Albinism Type 2 (OCA2). *Am J Hum Genet*. 73(3): 638-645.